

Hedda



Parentage

Öjebyn x Melahati

Breeding Center

?



Plant growth

medium

Plant Habit

semi-upright

Time of flowering

early

Time of Fruit Ripening

medium

Fruit Yield

medium

Susceptibility to main diseases and pests *

Powdery Mildew

1

Antracnose

5

WPBR

5

Gall Mite

5

* *Ranking scale: 1-9 (1 = resistant, 9 = very highly infected)*

Fruit Size / Weight

medium (0,71-1,10g)

Fruit Quality / Chemical Fruit Analyses *

soluble solids

medium **acidity** medium **anthocyanins** medium **ascorbic acids** very low

* *Ranking scale: very low, low, medium, high, very high*



Commercial use

Country specifics

other

Ben Nare



Parentage

Ben Lomond x ND 17/7

Breeding Center

James Hutton Institute, Dundee



Plant growth

tall

Plant Habit

upright

Time of flowering

early

Time of Fruit Ripening

early

Fruit Yield

medium

Susceptibility to main diseases and pests *

Powdery Mildew

1

Antracnose

5

WPBR

5

Gall Mite

5

** Ranking scale: 1-9 (1 = resistant, 9 = very highly infected)*

Fruit Size / Weight

medium (0,71-1,10g)

Fruit Quality / Chemical Fruit Analyses *

soluble solids

highacidityhighanthocyaninshighascorbic acidsmedium

** Ranking scale: very low, low, medium, high, very high*



Commercial use

Country specifics

other

Sunniva



Parentage

Öjebyn x Melahati

Breeding Center

?



Plant growth

medium

Plant Habit

semi-upright

Time of flowering

early

Time of Fruit Ripening

medium

Fruit Yield

medium

Susceptibility to main diseases and pests *

Powdery Mildew

1

Antracnose

5

WPBR

5

Gall Mite

5

** Ranking scale: 1-9 (1 = resistant, 9 = very highly infected)*

Fruit Size / Weight

small (<0,70g)

Fruit Quality / Chemical Fruit Analyses *

soluble solids

medium **acidity** medium **anthocyanins** low **ascorbic acids** medium

** Ranking scale: very low, low, medium, high, very high*



Commercial use
Country specifics
other

Narve Viking



Parentage

Ben Tron x SCRI C2/1/62

Breeding Center

?



Plant growth

tall

Plant Habit

upright

Time of flowering

late

Time of Fruit Ripening

late

Fruit Yield

high

Susceptibility to main diseases and pests *

Powdery Mildew

1

Antracnose

2

WPBR

5

Gall Mite

5

** Ranking scale: 1-9 (1 = resistant, 9 = very highly infected)*

Fruit Size / Weight

medium (0,71-1,10g)

Fruit Quality / Chemical Fruit Analyses *

soluble solids

highaciditymediumanthocyaninsmediumascorbic acidshigh

** Ranking scale: very low, low, medium, high, very high*



Commercial use

Country specifics

other

Ben Nevis



Parentage

(Brøtorp x Janslunda) x (Consort x Magnus)

Breeding Center

James Hutton Institute, Dundee



Plant growth

tall

Plant Habit

semi-upright

Time of flowering

early

Time of Fruit Ripening

early

Fruit Yield

high

Susceptibility to main diseases and pests *

Powdery Mildew

7

Antracnose

5

WPBR

5

Gall Mite

7

** Ranking scale: 1-9 (1 = resistant, 9 = very highly infected)*

Fruit Size / Weight

medium (0,71-1,10g)

Fruit Quality / Chemical Fruit Analyses *

soluble solids

highacidityhighanthocyaninshighascorbic acidshigh

* *Ranking scale: very low, low, medium, high, very high*



Commercial use

Country specifics

other

Tisel



Parentage

Titania x selfed

Breeding Center

Instytut Sadownictwa i Kwiaciarnictwa, Skierniewice



Plant growth

tall

Plant Habit

upright

Time of flowering

early

Time of Fruit Ripening

early

Fruit Yield

high

Susceptibility to main diseases and pests *

Powdery Mildew

2

Antracnose

5

WPBR

2

Gall Mite

2

** Ranking scale: 1-9 (1 = resistant, 9 = very highly infected)*

Fruit Size / Weight

medium (0,71-1,10g)

Fruit Quality / Chemical Fruit Analyses *

soluble solids

high**acidity**low**anthocyanins**low**ascorbic acids**very high

* *Ranking scale: very low, low, medium, high, very high*



Commercial use

Country specifics

other

Tiben



Parentage

Titania x Ben Nevis

Breeding Center

Instytut Sadownictwa i Kwiaciarnictwa, Skierniewice



Plant growth

tall

Plant Habit

semi-upright

Time of flowering

late

Time of Fruit Ripening

medium

Fruit Yield

medium

Susceptibility to main diseases and pests *

Powdery Mildew

2

Antracnose

5

WPBR

5

Gall Mite

5

** Ranking scale: 1-9 (1 = resistant, 9 = very highly infected)*

Fruit Size / Weight

medium (0,71-1,10g)

Fruit Quality / Chemical Fruit Analyses *

soluble solids

medium **acidity** high **anthocyanins** high **ascorbic acids** medium

* *Ranking scale: very low, low, medium, high, very high*



Commercial use

Country specifics

other

Tenah



Parentage

(Goliath x unknoww) x Brödtorp

Breeding Center

?



Plant growth

very tall

Plant Habit

upright

Time of flowering

early

Time of Fruit Ripening

early

Fruit Yield

high

Susceptibility to main diseases and pests *

Powdery Mildew

?

Antracnose

?

WPBR

?

Gall Mite

?

** Ranking scale: 1-9 (1 = resistant, 9 = very highly infected)*

Fruit Size / Weight

?

Fruit Quality / Chemical Fruit Analyses *

soluble solids

?acidity?anthocyanins?ascorbic acids?

* *Ranking scale: very low, low, medium, high, very high*



Commercial use

Country specifics

other

Ometa



Parentage

Westra x Seedling resistant to mildew

Breeding Center

?



Plant growth

tall

Plant Habit

upright

Time of flowering

late

Time of Fruit Ripening

medium

Fruit Yield

medium

Susceptibility to main diseases and pests *

Powdery Mildew

2

Antracnose

?

WPBR

?

Gall Mite

2

** Ranking scale: 1-9 (1 = resistant, 9 = very highly infected)*

Fruit Size / Weight

?

Fruit Quality / Chemical Fruit Analyses *

soluble solids

?acidity?anthocyanins?ascorbic acids?

* *Ranking scale: very low, low, medium, high, very high*



Commercial use

Country specifics

other

Kristin



Parentage

Ben Tron x L I 11/46-85 (Hedda x E.M.1428/70)

Breeding Center

?



Plant growth

tall

Plant Habit

upright

Time of flowering

early

Time of Fruit Ripening

early

Fruit Yield

medium

Susceptibility to main diseases and pests *

Powdery Mildew

1

Antracnose

5

WPBR

5

Gall Mite

5

** Ranking scale: 1-9 (1 = resistant, 9 = very highly infected)*

Fruit Size / Weight

large (1,11-1,50g)

Fruit Quality / Chemical Fruit Analyses *

soluble solids

medium**acidity**low**anthocyanins**medium**ascorbic acids**low

* *Ranking scale: very low, low, medium, high, very high*



Commercial use

Country specifics

other

Intercontinental



Parentage

Clone x Titania

Breeding Center

?

Sweden

Plant growth

tall

Plant Habit

semi-upright

Time of flowering

medium

Time of Fruit Ripening

medium

Fruit Yield

high

Susceptibility to main diseases and pests *

Powdery Mildew

5

Antracnose

5

WPBR

5

Gall Mite

5

** Ranking scale: 1-9 (1 = resistant, 9 = very highly infected)*

Fruit Size / Weight

very large (>1,50g)

Fruit Quality / Chemical Fruit Analyses *

soluble solids

?acidity?anthocyanins?ascorbic acids?

* *Ranking scale: very low, low, medium, high, very high*

?



Commercial use

Country specifics

other

Cheresnheva - Syn.Neva



Parentage

?

Breeding Center

Institute of Horticulture of the National Academy of Agrarian Sciences of Ukraine



Plant growth

medium

Plant Habit

upright

Time of flowering

early

Time of Fruit Ripening

early

Fruit Yield

high

Susceptibility to main diseases and pests *

Powdery Mildew

1

Antracnose

?

WPBR

?

Gall Mite

1

** Ranking scale: 1-9 (1 = resistant, 9 = very highly infected)*

Fruit Size / Weight

large (1,11-1,50g)

Fruit Quality / Chemical Fruit Analyses *

soluble solids

?**acidity**low**anthocyanins**?**ascorbic acids**?

* *Ranking scale: very low, low, medium, high, very high*



Commercial use

Country specifics

other